

Connecting via Winsock to STN

Trying 3106016892...Open

Welcome to STN International! Enter x:x
LOGINID:sssptal644axd
PASSWORD:
TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files
NEWS 3 Feb 06 Engineering Information Encompass files have new names
NEWS 4 Feb 16 TOXLINE no longer being updated
NEWS 5 Apr 23 Search Derwent WPINDEK by chemical structure
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA

NEWS EXPRESS April 18 CURRENT WINDOWS VERSION IS V6.0,
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),
AND CURRENT DISCOVER FILE IS DATED 04/06
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 09:09:51 ON 30 APR 2001

=> file medline caplus embase biosis
COST IN U.S. DOLLARS SINCE FILE ENTRY TOTAL SESSION
FULL ESTIMATED COST 0.15 0.15

FILE 'MEDLINE' ENTERED AT 09:10:09 ON 30 APR 2001

FILE 'CAPLUS' ENTERED AT 09:10:09 ON 30 APR 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 09:10:09 ON 30 APR 2001
COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 09:10:09 ON 30 APR 2001
COPYRIGHT (C) 2001 BIOSIS(R)

=> s MIIC and A20
L1 11 MIIC AND A20

=> dup rem l1
PROCESSING COMPLETED FOR L1
L2 3 DUP REM L1 (8 DUPLICATES REMOVED)

=> s l2 (P) class
PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH
FIELD CODE - 'AND' OPERATOR ASSUMED 'L7 (P) CLASS'
PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH
FIELD CODE - 'AND' OPERATOR ASSUMED 'L9 (P) CLASS'
PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH
FIELD CODE - 'AND' OPERATOR ASSUMED 'L11 (P) CLASS'
PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH
FIELD CODE - 'AND' OPERATOR ASSUMED 'L13 (P) CLASS'
L3 3 L2 (P) CLASS

=> dis l3 1-3 kwic ibib abs

L3 ANSWER 1 OF 3 MEDLINE
TI Major histocompatibility complex **class II** compartments in human and mouse B lymphoblasts represent conventional endocytic compartments.
AB In most human and mouse antigen-presenting cells, the majority of intracellular major histocompatibility complex (MHC) **class II** molecules resides in late endocytic MHC **class II** compartments (MIICs), thought to function in antigen processing and peptide loading. However, in mouse A20 B cells, early endocytic **class II**-containing vesicles (CIIVs) have been reported to contain most of the intracellular MHC **class II** molecules and have also been implicated in formation of MHC **class II**-peptide complexes. To address this discrepancy, we have studied in great detail the endocytic pathways of both a human (6H5.DM) and a mouse (A20.Ab) B cell line. Using quantitative immunoelectron microscopy on cryosections of cells that had been pulse-chased with transferrin-HRP or BSA-gold as endocytic tracers, we have identified up to six endocytic subcompartments including an early MIIC type enriched in invariant chain, suggesting that it serves as an important entrance to the endocytic pathway for newly synthesized MHC **class II**/invariant chain complexes. In addition, early MIICs represented the earliest endocytic compartment containing MHC **class II**-peptide complexes, as shown by using an antibody against an abundant endogenous **class II**-peptide complex. The early MIIC exhibited several though not all of the characteristics reported for the CIIV and was situated just downstream of early endosomes. We have not encountered any special **class II**-containing endocytic structures besides those normally present in nonantigen-presenting cells. Our results therefore suggest that B cells use conventional endocytic compartments rather than having developed a unique compartment to accomplish MHC **class II** presentation.

CT
Compartmentation: IM, immunology
Cell Line

Cell Line, Transformed
*Endocytosis
Endocytosis: IM, immunology
Gold Colloid: ME, metabolism
HLA-D Antigens: ME, metabolism
Histocompatibility Antigens Class II: IM, immunology
Histocompatibility Antigens Class II: ME, metabolism
*Histocompatibility Antigens Class II: PH, physiology
Horseradish Peroxidase: ME, metabolism
Kinetics
Lymphocyte Transformation
Mice
Serum Albumin, Bovine: ME, metabolism
Transferrin: ME, . . .

CN. . . (Antibodies, Monoclonal); 0 (Antigens, Differentiation, B-Lymphocyte); 0 (Gold Colloid); 0 (H2-M antigens); 0 (HLA-D Antigens); 0 (HLA-DMB); 0 (Histocompatibility Antigens Class II); 0 (Serum Albumin, Bovine); 0 (invariant chain); EC 1.11.1.- (Horseradish Peroxidase)

ACCESSION NUMBER: 1998012197 MEDLINE
DOCUMENT NUMBER: 98012197 PubMed ID: 9348281
TITLE: Major histocompatibility complex class II compartments in human and mouse B lymphoblasts represent conventional endocytic compartments.
AUTHOR: Kleijmeer M J; Morkowski S; Griffith J M; Rudensky A Y; Geuze H J
CORPORATE SOURCE: Department of Cell Biology, School of Medicine and Institute of Biomembranes, Utrecht University, 3584 CX Utrecht, The Netherlands.
SOURCE: JOURNAL OF CELL BIOLOGY, (1997 Nov 3) 139 (3) 639-49. Journal code: HMV; 0375356. ISSN: 0021-9525.
PUB. COUNTRY: United States
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199711
ENTRY DATE: Entered STN: 19971224
Last Updated on STN: 19971224
Entered Medline: 19971121

AB In most human and mouse antigen-presenting cells, the majority of intracellular major histocompatibility complex (MHC) class II molecules resides in late endocytic MHC class II compartments (MIICs), thought to function in antigen processing and peptide loading. However, in mouse A20 B cells, early endocytic class II-containing vesicles (CIIVs) have been reported to contain most of the intracellular MHC class II molecules and have also been implicated in formation of MHC class II-peptide complexes. To address this discrepancy, we have studied in great detail the endocytic pathways of both a human (6H5.DM) and a mouse (A20.Ab) B cell line. Using quantitative immunoelectron microscopy on cryosections of cells that had been pulse-chased with transferrin-HRP or BSA-gold as endocytic tracers, we have identified up to six endocytic subcompartments including an early MIIC type enriched in invariant chain, suggesting that it serves as an important entrance to the endocytic pathway for newly synthesized MHC class II/invariant chain complexes. In addition, early MIICs represented the earliest endocytic compartment containing MHC class II-peptide complexes, as shown by using an antibody against an abundant endogenous class II-peptide complex. The early MIIC exhibited several though not all of the characteristics reported for the CIIV and was situated just downstream of early endosomes. We have not encountered any special class II-containing endocytic structures besides those normally present in nonantigen-presenting cells. Our results therefore suggest that B cells use conventional endocytic compartments rather than having developed a unique compartment to accomplish MHC class II presentation.

L3 ANSWER 2 OF 3 MEDLINE
TI Ii chain controls the transport of major histocompatibility complex class II molecules to and from lysosomes.

AB Major histocompatibility complex class II molecules are synthesized as a nonameric complex consisting of three alpha beta dimers associated with a trimer of invariant chain. . . in the Ii chain cytoplasmic domain directs the complex to endosomes where Ii chain is proteolytically processed and removed, allowing class II molecules to bind antigenic peptides before reaching the cell surface. Ii chain dissociation and peptide binding are thought to occur in one or more postendosomal sites related either to endosomes (designated CIIV) or to lysosomes (designated MIIC). We now find that in addition to initially targeting alpha beta dimers to endosomes, Ii chain regulates the subsequent transport of class II molecules. Under normal conditions, murine A20 B cells transport all of their newly synthesized class II I-A(b) alpha beta dimers to the plasma membrane with little if any reaching lysosomal compartments. Inhibition of Ii processing. . . cysteine/serine protease inhibitor leupeptin, however, blocked transport to the cell surface and caused a dramatic but selective accumulation of I-A(b) class II molecules in lysosomes. In leupeptin, I-A(b) dimers formed stable complexes with a 10-kD NH2-terminal Ii chain fragment (Ii-p10), normally. . . Our results suggest that alterations in the rate or efficiency of Ii chain processing can alter the postendosomal sorting of class II molecules, resulting in the increased accumulation of alpha beta dimers in lysosome-like MIIC. Thus, simple differences in Ii chain processing may account for the highly variable amounts of class II found in lysosomal compartments of different cell types or at different developmental stages.

CT . . .
Cell Compartmentation: PH, physiology
Cell Membrane: CH, chemistry
Cell Membrane: ME, metabolism
Cell Membrane: UL, ultrastructure
Dimerization
Electrophoresis: MT, methods
Histocompatibility Antigens Class II: CH, chemistry
Histocompatibility Antigens Class II: DE, drug effects
*Histocompatibility Antigens Class II: ME, metabolism
Leupeptins: PD, pharmacology
Lymphoma
Lysosomes: CH, chemistry
*Lysosomes: ME, metabolism
Lysosomes: UL, ultrastructure
Mice
Microscopy, . . .

CN 0 (Antigens, Surface); 0 (Histocompatib Antigens Class II);
 0 (Leupeptins)
 ACCESSION NUMBER: 97258865 MEDLINE
 DOCUMENT NUMBER: 97258865 PubMed ID: 9105036
 TITLE: Ii chain controls the transport of major histocompatibility complex class II molecules to and from lysosomes.
 AUTHOR: Brachet V; Raposo G; Amigorena S; Mellman I
 CORPORATE SOURCE: Institut Curie, Section de Recherche Institut National de la Sante et de la Recherche Medicale C/JF-95.01, Paris, France.
 SOURCE: JOURNAL OF CELL BIOLOGY, (1997 Apr 7) 137 (1) 51-65.
 Journal code: HMV; 0375356. ISSN: 0021-9525.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199705
 ENTRY DATE: Entered STN: 19970514
 Last Updated on STN: 19970514
 Entered Medline: 19970508

AB Major histocompatibility complex class II molecules are synthesized as a nonameric complex consisting of three alpha beta dimers associated with a trimer of invariant (Ii) chains. After exiting the TGN, a targeting signal in the Ii chain cytoplasmic domain directs the complex to endosomes where Ii chain is proteolytically processed and removed, allowing class II molecules to bind antigenic peptides before reaching the cell surface. Ii chain dissociation and peptide binding are thought to occur in one or more postendosomal sites related either to endosomes (designated CIIV) or to lysosomes (designated MIIC). We now find that in addition to initially targeting alpha-beta dimers to endosomes, Ii chain regulates the subsequent transport of class II molecules. Under normal conditions, murine A20 B cells transport all of their newly synthesized class II I-A(b) alpha beta dimers to the plasma membrane with little if any reaching lysosomal compartments. Inhibition of Ii processing by the cysteine/serine protease inhibitor leupeptin, however, blocked transport to the cell surface and caused a dramatic but selective accumulation of I-A(b) class II molecules in lysosomes. In leupeptin, I-A(b) dimers formed stable complexes with a 10-kD NH2-terminal Ii chain fragment (Ii-p10), normally a transient intermediate in Ii chain processing. Upon removal of leupeptin, Ii-p10 was degraded and released, I-A(b) dimers bound antigenic peptides, and the peptide-loaded dimers were transported slowly from lysosomes to the plasma membrane. Our results suggest that alterations in the rate or efficiency of Ii chain processing can alter the postendosomal sorting of class II molecules, resulting in the increased accumulation of alpha beta dimers in lysosome-like MIIC. Thus, simple differences in Ii chain processing may account for the highly variable amounts of class II found in lysosomal compartments of different cell types or at different developmental stages.

L3 ANSWER 3 OF 3 MEDLINE
 TI HLA-DM is localized to conventional and unconventional MHC class II-containing endocytic compartments.
 AB HLA-DM molecules remove invariant (Ii) chain peptides from newly synthesized MHC class II complexes. Their localization may thus delineate compartments, e.g., MIIC, specialized for loading peptides onto class II molecules. In murine A20 B cells, however, DM is not restricted to specialized endosomal class II-containing vesicles (CIIV). Although DM was found in CIIV, it was also found throughout the endocytic pathway, principally in lysosomes devoid of class II molecules. In human lymphoblasts, HLA-DM was found in structures indistinguishable from late endosomes or lysosomes, although in these cells the lysosomes contained MHC class II molecules. Thus, the distribution of HLA-DM does not necessarily identify specialized class II compartments. Many "MIIC" may represent conventional lysosomes that accumulate MHC class II and HLA-DM in a number of cell types.

ACCESSION NUMBER: 96209673 MEDLINE
 DOCUMENT NUMBER: 96209673 PubMed ID: 8624813
 TITLE: HLA-DM is localized to conventional and unconventional MHC class II-containing endocytic compartments.
 AUTHOR: Pierre P; Denzin L K; Hammond C; Drake J R; Amigorena S; Cresswell P; Mellman I
 CORPORATE SOURCE: Department of Cell Biology, Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, Connecticut 06520-8002, USA.
 SOURCE: IMMUNITY, (1996 Mar) 4 (3) 229-39.
 Journal code: CCF; 9432918. ISSN: 1074-7613.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199606
 ENTRY DATE: Entered STN: 19960708
 Last Updated on STN: 19960708
 Entered Medline: 19960627

AB HLA-DM molecules remove invariant (Ii) chain peptides from newly synthesized MHC class II complexes. Their localization may thus delineate compartments, e.g., MIIC, specialized for loading peptides onto class II molecules. In murine A20 B cells, however, DM is not restricted to specialized endosomal class II-containing vesicles (CIIV). Although DM was found in CIIV, it was also found throughout the endocytic pathway, principally in lysosomes devoid of class II molecules. In human lymphoblasts, HLA-DM was found in structures indistinguishable from late endosomes or lysosomes, although in these cells the lysosomes contained MHC class II molecules. Thus, the distribution of HLA-DM does not necessarily identify specialized class II compartments. Many "MIIC" may represent conventional lysosomes that accumulate MHC class II and HLA-DM in a number of cell types.

=> end

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:y
 COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
7.59	7.74

FULL ESTIMATED COST

STN INTERNATIONAL LOGOFF AT 09:11:41 ON 30 APR 2001